

Nobel Prize in Medicines 1944



Joseph Erlanger



Herbert Spencer Gasser

The Nobel Prize in Physiology or Medicine 1944 was awarded jointly to Joseph Erlanger and Herbert Spencer Gasser "for their discoveries relating to the highly differentiated functions of single nerve fibres"

Three great electrophysiological discoveries can be regarded as milestones in the development of our knowledge of nerve physiology. In the middle of the last century, long before Alfred Nobel had bequeathed this great fund to the world, Du Bois-Reymond showed that the nerve impulse was an electrical wave of negativity transmitted along the nerve. Helmholtz made the first measurements of its average speed of propagation in a nerve stem. The second great discovery, rewarded with a share in the Nobel Prize for 1932, was Adrian's observation that sense organs and nerve cells discharge whole series of such impulses. In each individual fibre the nerve impulse is of constant size but the stronger the stimulus, the greater the frequency of the impulses discharged along the nerve. The nerve cells, as it were, communicate with each other by a kind of machine-gun fire. This description of the mechanism is inadequate from the standpoint of physics but otherwise illuminating. And, indeed, if the impulses are led to a loudspeaker over an amplifier it does

sound like machine-gun fire. The discoveries of Erlanger and Gasser constitute a third step forwards.

In 1907 the Swedish physiologist Gustaf Göthlin made the assumption that conduction velocity in thick nerve fibres is greater than in thin ones. The basis for this view was W. Thompson's formula for electrical-cable conduction. This assumption gave a physiological interpretation of the well-known fact that the individual fibres of a nerve stem vary in cross section. Some fibres are less than 0.001 mm in diameter, others just above 0.020 mm. Lopicque and his colleagues, from 1913 onwards, published some papers in which indirect evidence in support of this view was advanced. In a series of remarkable papers - remarkable in respect of both technique and wealth of new information unearthed - Erlanger and Gasser proved this hypothesis to have been correct. As so often happens in experimental sciences, the additional steps necessary for full clarity as well as the development of an elegant new technique, heralded an experimental expansion of great width and significance. The seemingly simple cables turned out to have been endowed with a high degree of differentiation. Since nerve fibres are to be regarded as extensions of nerve cells these results are indeed of extreme importance for the physiology of the higher centres such as the brain and the spinal cord. This fact should be given especial consideration in appraising the significance of the work of Erlanger and Gasser.

Erlanger and Gasser showed that the nerve fibres, according to their conduction velocities, could be divided into three main groups of which the first, group A, could be further subdivided. The thickest mammalian fibres, the A-fibres, conduct impulses as fast as from 5 to 100 metre per second, the thinnest, the C-fibres, have conduction velocities below 2 metre per second. Between these two groups there are the B-fibres with conduction velocities from 3 to 14 metre per second. A large number of other properties of the nerve fibres vary with the speed of conduction, for instance, the duration of the impulse, its rate of rise, its size, the duration of the inexcitable or refractory period following each impulse, the threshold of excitation, the sensitivity of the discharge to pressure on the nerve and to asphyxia, in short, an array of properties connected with

impulse conduction all of which need not vary in an exactly parallel manner. Erlanger and Gasser also showed how this highly differentiated system with its three main types of fibre was distributed over the in- and outgoing fibres of the spinal cord, the so-called sensory and motor roots. The perception of pain is largely mediated by very thin, slowly conducting fibres, muscle sense and touch by rapidly conducting fibres. The muscles of the body are also thrown into movement by fast fibres.

In the brain and the spinal cord the time ratios of the impulses are of primary importance for the cooperation of the nerve cells. A difference of 0.001-0.005 seconds in the time of arrival of impulses means that a given path may be found opened or closed for their passage onwards. Problems of this kind belong to the present-day programme of experimentation in these fields.

The admirable technique of Erlanger and Gasser soon showed them a road to new discoveries, chiefly concerned with the changes of excitability that occur at a nerve cross section at which impulses arrive. The arrival of one or several impulses to such a region was found to be followed by slow changes of excitability which were associated with slow changes of electrical potential, studied in detail by Gasser. These changes of excitability enhance or depress succeeding impulses. Such «after-potentials» had been seen before, but Gasser and his collaborators demonstrated their independent character and showed that they behaved in a different manner in the three main types of fibre. The concept of a high degree of differentiation of the nerve fibres for their different tasks was thus again supported by a new group of facts. These are of particular importance for the physiology of the central nervous system. A prominent feature of this region is interaction between excitation and inhibition in close association with slow potential changes. Erlanger and his collaborators devoted themselves to an analysis of the changes of excitability in a nerve influenced by a constant electrical current. One of their most important discoveries was the demonstration that sensory nerves in many respects differed from motor nerves. The sensory nerves had, for instance, lower thresholds of excitation and they put up less

resistance to impulse generation (less «accommodation») than motor nerves. This fresh contribution to the differentiation among the nerve fibres has far-reaching consequences.

When today Erlanger and Gasser receive the 1944 Nobel Prize for their discoveries concerning the highly differentiated properties of single nerve fibres, it might be pointed out that their achievement was not born, fixed and armoured, in the manner of the birth of Pallas Athene. But no sooner had their first result given them the key word than discovery followed hard upon discovery until their colleagues everywhere in the world came to realize that a great new synthesis had been born to nerve physiology. This synthesis is based on new facts, well-hardened by a masterly technique cementing them into a groundwork on which will be erected whatever structure the future has in store for the physiology of the central and peripheral nervous system.

For more details please visit:

http://www.nobelprize.org/nobel_prizes/medicine/laureates/1944/press.html